

# Stability of the entropy for superstatistics

A. M. C. Souza<sup>1,2</sup> and C. Tsallis<sup>1</sup> \*

<sup>1</sup>*Centro Brasileiro de Pesquisas Fisicas*

*Rua Xavier*

*Sigaud 150, 22290-180 Rio de Janeiro-RJ, Brazil*

<sup>2</sup>*Departamento de Fisica*

*Universidade Federal de*

*Sergipe, 49100-000 Sao Cristovao-SE, Brazil*

## Abstract

The Boltzmann-Gibbs celebrated entropy  $S_{BG} = -k \sum_i p_i \ln p_i$  is *concave* (with regard to all probability distributions  $\{p_i\}$ ) and *stable* (under arbitrarily small deformations of any given probability distribution). It seems reasonable to consider these two properties as *necessary* for an entropic form to be a *physical* one in the thermostatistical sense. Most known entropic forms (e.g., Renyi entropy) violate these conditions, in contrast with the basis of nonextensive statistical mechanics, namely  $S_q = k \frac{1 - \sum_i p_i^q}{q-1}$  ( $q \in \mathcal{R}$ ;  $S_1 = S_{BG}$ ), which satisfies both ( $\forall q > 0$ ). We have recently generalized  $S_q$  (into  $S$ ) in order to yield, through optimization, the Beck-Cohen superstatistics. We show here that  $S$  satisfies both conditions as well. Given the fact that the (experimentally observed) optimizing distributions are invariant through *any* monotonic function of the entropic form to be optimized, this might constitute a very strong criterion for identifying the physically correct entropy.

---

\*amcsouza@ufs.br, tsallis@cbpf.br

The Boltzmann-Gibbs (BG) entropy expression

$$S_{BG} = -k \sum_{i=1}^W p_i \ln p_i, \quad (1)$$

(sometimes referred to as the *Boltzmann-Gibbs-Shannon entropy*), and its equal-probability particular case  $S_{BG} = k \ln W$ , undoubtedly constitute a major step in human understanding of the underlying laws of nature. Its continuous form  $S_{BG} = -k \int dx p(x) \ln p(x)$ , historically introduced by Boltzmann and Gibbs, its quantum form  $S_{BG} = -k \text{Tr } \rho \ln \rho$ , introduced by von Neumann, and the above mentioned discrete form (1), introduced by Shannon (though in a different context), have been at the basis of invaluable scientific results in physics, chemistry and elsewhere. Let us learn more from it. The optimization of Eq. (1) under the canonical ensemble (equilibrium with a thermostat) constraints  $\sum_{i=1}^W p_i = 1$  and  $\sum_{i=1}^W p_i E_i = U_{BG}$  ( $U_{BG}$  is the internal energy, and  $\{E_i\}$  are the eigenvalues of the Hamiltonian), yields the celebrated BG probability for the equilibrium stationary state

$$p_i \propto e^{-\beta E_i} \quad (\beta \equiv 1/kT) \quad (2)$$

This distribution can always be, at least in principle, experimentally checked (and it has been so with well known success). Not so  $S_{BG}$ ! Although it normally goes without discussion, it is important to notice, for the arguments we want to develop here, that *any* monotonic function of  $S_{BG}$  (e.g.,  $S_{BG}^3/k^2$ ) would provide, through optimization, the very same distribution as Eq. (2). BG statistical mechanics is however much more than just the equilibrium distribution, and there are plethora reasons for considering the *physical entropy* to be precisely  $S_{BG}$  and no other function of it. We will quickly realize that the situation can become much less trivial as soon as we want to generalize BG statistical mechanics, in order to address other (typically nonergodic) stationary states such as those which ubiquitously emerge in complex systems.

In order to handle nonequilibrium states of large classes of systems (examples nowadays known include turbulence [1–3], electron-positron annihilation [4], quark-gluon plasma [5], anomalously diffusing micro-organisms [6], classical [7] and quantum [8] chaos, long-range

many-body Hamiltonians [9], economics [10], and others; see [11] for reviews), one of us postulated [12] the entropic form

$$S_q = k \frac{1 - \sum_{i=1}^W p_i^q}{q - 1} \quad (q \in \mathcal{R}; S_1 = S_{BG}), \quad (3)$$

as the basis for generalizing BG statistical mechanics. It can be trivially verified that  $p_{ij}^{A+B} = p_i^A \times p_j^B$ ,  $\forall (i, j)$ , implies  $S_q(A+B)/k = S_q(A)/k + S_q(B)/k + (1-q)S_q(A)S_q(B)/k^2$ , property which makes the generalized formalism to be sometimes referred to as *nonextensive statistical mechanics*. If we optimize  $S_q$  with the already mentioned norm constraint and  $\sum_{i=1}^W p_i^q E_i / \sum_{i=1}^W p_i^q = U_q$ , we obtain [13]

$$p_i \propto e_q^{-\beta'_q E_i}, \quad (4)$$

where the  $q$ -exponential function is defined as  $e_q^x \equiv [1 + (1-q)x]^{1/(1-q)}$  ( $e_1^x = e^x$ ), and  $\beta'_q$  is an effective inverse temperature. This distribution is ubiquitously detected in nature. The question arises: *is  $S_q$  the proper physical entropy for generalizing the BG formalism?* Indeed, any monotonic function of  $S_q$ , such as for example the Renyi entropy  $S_q^R \equiv [\ln \sum_{i=1}^W p_i^q] / (1-q) = (1-q)^{-1} \ln[1 + (1-q)S_q]$  (we use  $k = 1$  from now on), will also yield Eq. (4) if optimized under the same constraints. The answer seems to be *yes*. Indeed,  $S_q$  is,  $\forall q > 0$ , both *concave* (with regard to all probability distributions  $\{p_i\}$ ) and *stable* (under arbitrarily small deformations of any given probability distribution), whereas  $S_q^R$  is *not* (like all other known entropic forms which are monotonic functions of  $S_q$ ) [14,15].

The entropy  $S_{BG}$  can be obtained from appropriate set of axioms, as long shown by Shannon and also by Khinchin. However, the same occurs with  $S_q$  and  $S_q^R$  (and others) by modifying in admissible manners Shannon's and/or Khinchin's sets of axioms [16]. Therefore this path does not help much for distinguishing the physical entropy from its monotonic functions. But concavity and stability do. Indeed, no monotonic function of  $S_q$  is known which satisfies both. On top of this,  $S_q$  allows, in all known cases (both vanishing and nonvanishing Lyapunov exponents, i.e., both zero and positive Kolmogorov-Sinai entropy), for *finite* entropy production per unit time [7], whereas the others do not.

Now, Wilk and Włodarczyk [17] as well as Beck [2], noticed that Eq. (4) can be seen as the result of complex dynamics in stationary states associated with regions that exhibit spatio-temporal fluctuations of temperature. Such interpretation relies on the fact that  $e_q^{-\beta'_q E_i} = \int_0^\infty d\beta e^{-\beta E_i} f_q(\beta)$ , where  $f_q(\beta)$  is a  $\chi^2$  distribution, the entropic index  $q$  being given by  $q = \langle \beta^2 \rangle / \langle \beta \rangle^2$ . Clearly,  $\lim_{q \rightarrow 1} f_q(\beta) = \delta(\beta - \beta'_1)$ , where  $\delta$  represents Dirac's delta.

Very recently, Beck and Cohen [18] made a further step along this line, which they called *superstatistics* (“statistics of statistics”). They introduced a generalized BG factor, namely

$$B(E) = \int_0^\infty d\beta e^{-\beta E} f(\beta), \quad (5)$$

where  $f(\beta)$  is a generic admissible probability distribution of the stochastic inverse “temperature” satisfying  $\int_0^\infty d\beta f(\beta) = 1$ , and

$$q_{BC} \equiv \langle \beta^2 \rangle / \langle \beta \rangle^2, \quad (6)$$

where  $\langle \dots \rangle \equiv \int_0^\infty d\beta (\dots) f(\beta)$ . We have slightly modified their original notation, by adding the subindex  $BC$  (which stands for “Beck-Cohen”), in order to avoid confusion with  $q$ . If  $f(\beta)$  is a  $\chi^2$  (or gamma) distribution, then  $q_{BC} = q$ ; otherwise, those are generically different quantities.

The reason for considering the generalization (5) of the  $q$ -exponential weight appearing in nonextensive statistical mechanics basically is that, for systems with sufficiently complex dynamics, there is no *a priori* reason for not expecting the existence in nature of even more general distributions. The effective statistical mechanics of such systems will depend on the statistical properties of the fluctuations of the temperature, or of even other intensive quantities. Naturally, if there are no fluctuations of intensive quantities at all, the system must obey BG statistical mechanics (i.e.,  $q_{BC} = q = 1$ ). Beck and Cohen also showed that, for small variance of the fluctuations (i.e.,  $|q_{BC} - 1| \ll 1$ ), the first order correction to the BG statistics is the nonextensive statistics [12]. That is, the particular mechanism where nonextensivity is driven by relative small fluctuations of temperature is *universal* and precisely corresponds to nonextensive statistical mechanics with  $q = q_{BC}$ . The whole idea was also illustrated with the uniform, bimodal, log-normal and  $F$ -distributions.

So far so good. But it is important to realize that having a generalized statistical weight such as (5) is *not* enough for having a statistical mechanics. Indeed, consistent expressions for the entropy  $S$  and for the internal energy  $U$  are also needed. By “consistent” we mean that the optimization of  $S$  with the norm and the energy constraints should yield Eq. (5). We have recently bridged this gap [19] and proposed the entropic functional  $S$ , as well as the associated energy constraint, so that superstatistics can be now *derived* from a variational principle within a statistical mechanical frame. Let us recall that any monotonic function of the entropic functional  $S$  introduced in [19] will also lead to Eq. (5). The entropy  $S$  seems, however, to be the *physical* one. Indeed, not only it is concave by construction [19], but it is also stable. The proof of the latter is the main result of the present paper.

The entropic form  $S$  and the associated constraint for the internal energy  $U$  introduced in [19] are given by (from now on  $k = 1$  for simplicity)

$$S = \sum_{i=1}^W s(p_i) \quad (s(p_i) \geq 0; s(0) = s(1) = 0) , \quad (7)$$

and

$$\frac{\sum_{i=1}^W u(p_i) E_i}{\sum_{i=1}^W u(p_i)} = U \quad (0 \leq u(p_i) \leq 1; u(0) = 0; u(1) = 1) . \quad (8)$$

For example, for the BG statistics, we have  $s_{BG}(p) = -p \ln p$  and  $u_{BG}(p) = p$ , and, for the nonextensive statistics, we have  $s_q(p) = (p - p^q)/(q - 1)$  and  $u_q(p) = p^q$ . The function  $s(p)$  should generically have a definite concavity  $\forall p \in [0, 1]$ . Conditions (7) imply that  $S \geq 0$  and that certainty corresponds to  $S = 0$ . The function  $u(p)$  should generically be a monotonically increasing one. Certainty about  $E_j$  implies  $U = E_j$ . The quantity  $u(p_i)/\sum_{j=1}^W u(p_j)$  constitutes itself a probability distribution (the *escort distribution* [20]). Starting from the distribution of  $f(\beta)$ , we calculate  $B(E)$  and  $\int_0^\infty dE' B(E')$ , and we find the inverse function, noted  $E(p)$ , of  $p(E) \equiv B(E)/\int_0^\infty dE' B(E')$ . Now, two cases are possible. The first one corresponds to distributions where the lowest admissible value of  $E(p)$  is  $E^* \rightarrow -\infty$ . In this case [19],  $u(p) = p$  and

$$s(p) = \alpha p + \int_0^p dx E(x) , \quad (9)$$

where  $\alpha = -\int_0^1 dx E(x)$ . The second case corresponds to a finite and known value of  $E^*$ . In this case, the final form of  $s(p)$  is given by [19]

$$s(p) = E^*[p + u(p)], \quad (10)$$

where

$$u(p) = \frac{\int_0^p \frac{dx}{1-E(x)/E^*}}{\int_0^1 \frac{dx}{1-E(x)/E^*}}. \quad (11)$$

Eqs. (9), (10) and (11) completely determine the formulation of the statistical mechanics associated with the Beck-Cohen superstatistics.

For a statistical quantity  $O(p)$  to be an observable, or physical quantity, a necessary condition [14] is that, under arbitrary small variations of the probabilities  $p$ , its relative variation remains small. We then say that  $O(p)$  is *stable*. To be more precise, if we consider two probability sets  $p$  and  $p'$  associated with  $W$  microstates, the measure of the size of the deformation can be defined as follows [14]

$$||p - p'|| = \sum_{i=1}^W |p_i - p'_i|. \quad (12)$$

The condition of stability of  $O(p)$  is then given by

$$||p - p'|| < \delta_\epsilon \quad \Rightarrow \quad \left| \frac{O(p) - O(p')}{O_{\max}} \right| < \epsilon \quad (13)$$

for any  $\epsilon > 0$ , with  $\delta_\epsilon > 0$  being independent from  $W$ . This implies, in particular, that  $\lim_{\epsilon \rightarrow 0} \lim_{W \rightarrow \infty} R = \lim_{W \rightarrow \infty} \lim_{\epsilon \rightarrow 0} R = 0$ .

Using conditions (12) and (13), Lesche [14] showed that the Rényi entropies are unstable, and, therefore, cannot represent experimentally observable quantities (except, of course, for  $q = 1$ , when they recover the BG entropy, which is stable). Recently, using this formalism, Abe proved [21] that  $S_q$  is stable and can consequently provide the physical entropic basis for the power-law distributions frequently observed in nature (Abe also pointed that another candidate, namely the so called normalized nonextensive entropies, are unstable, like the Renyi ones). It is along these lines that we shall now prove that  $S$ , as given by Eqs. (9), (10) and (11), is stable.

We first remark that  $B(E)$  is a physical generalization of the  $q$ -exponential factor (hence of the BG factor), and must therefore be a monotonically decreasing function; the same holds then for  $p(E)$  and for  $E(p)$ . It immediately follows from Eq. (9) that

$$S(p) - S(p') = \sum_{i=1}^W \int_{p'_i}^{p_i} dx E(x) . \quad (14)$$

Notice that, since  $S(p)$  is concave, we can write  $S(\lambda p + (1 - \lambda)p') \geq \lambda S(p) + (1 - \lambda)S(p')$  ( $\forall 0 \leq \lambda \leq 1$ ). Since  $E(p)$  is positive and a monotonically decreasing function, it follows from Eq. (9) that  $F(p) \equiv \int_0^p dx E(x)$  is a concave monotonically increasing function. Using the monotonicity of  $E(x)$ , if  $p_i \geq p'_i \geq 0$ , we have that

$$\int_{p'_i}^{p_i} dx E(x) = \int_0^{p_i - p'_i} dx E(x + p'_i) \leq \int_0^{p_i - p'_i} dx E(x) = F(p_i - p'_i), \quad (15)$$

and, if  $p'_i \geq p_i \geq 0$ , we then have that

$$\int_{p'_i}^{p_i} dx E(x) \leq - \int_0^{p'_i - p_i} dx E(x) = -F(p'_i - p_i). \quad (16)$$

Considering the upperbound, we can write that

$$|S(p) - S(p')| \leq \sum_{i=1}^W \left| \int_{p'_i}^{p_i} dx E(x) \right| \leq \sum_{i=1}^W F(|p_i - p'_i|). \quad (17)$$

On the other hand, from (10) we obtain

$$S(p) - S(p') = E^* \sum_{i=1}^W [u(p_i) - u(p'_i)] . \quad (18)$$

Using that  $u(p)$  is a monotonically increasing function and that  $S(p)$  is concave, we can see from Eq. (10) that  $u(p)$  is a concave monotonically increasing function if  $E^* > 0$ , and it is a convex monotonically increasing function if  $E^* < 0$ . Hence,  $F'(p) = E^* u(p)$  always is a concave monotonically increasing function. It follows that, if  $p_i \geq p'_i \geq 0$ , we have  $F'(p_i) - F'(p'_i) \leq F'(p_i - p'_i)$ , and, if  $p'_i \geq p_i \geq 0$ , we have  $-[F'(p_i) - F'(p'_i)] \leq F'(p'_i - p_i)$ . This implies that  $|F'(p_i) - F'(p'_i)| \leq F'(|p_i - p'_i|)$ , then

$$|S(p) - S(p')| \leq \sum_{i=1}^W |F'(p_i) - F'(p'_i)| \leq \sum_{i=1}^W F'(|p_i - p'_i|) . \quad (19)$$

We now use that  $g(\lambda x)g(y) \leq g(x)g(\lambda y)$  for any increasing concave function  $g(x)$ , with  $g(0) \geq 0$ ,  $0 < x < y$  and  $0 \leq \lambda < 1$ . Considering  $\lambda = \|p - p'\| < 1$ ,  $x = \frac{|p_i - p'_i|}{\|p - p'\|}$  and  $y = 1$  [22], it follows that

$$g(|p_i - p'_i|) \leq \frac{g(\|p - p'\|)}{g(1)} g\left(\frac{|p_i - p'_i|}{\|p - p'\|}\right). \quad (20)$$

Finally, since  $F(p)$  and  $F'(p)$  are increasing concave functions, from Eqs.(17) and (19), if we identify  $g(p)$  as being  $F(p)$  or  $F'(p)$ , we have

$$|S(p) - S(p')| \leq \frac{g(\|p - p'\|)}{g(1)} \left[ S\left(\frac{|p_i - p'_i|}{\|p - p'\|}\right) + g(1) \right]. \quad (21)$$

Therefore, denoting by  $S_{\max}$  the maximal entropy, we obtain

$$|S(p) - S(p')| \leq \frac{g(\|p - p'\|)}{g(1)} [S_{\max} + g(1)]. \quad (22)$$

This inequality shows that, for  $\epsilon > 0$  such that  $|S(p) - S(p')| \leq \epsilon S_{\max}$ , we have that  $\|p - p'\| \leq \delta_\epsilon$ , which proves the stability of the entropy for superstatistics.

Summarizing, the entropies introduced in Ref. [19] have been studied. Since they are simultaneously concave and stable, it is allowed to think that these entropies are physical ones, and as such can be used to formulate a proper statistical mechanics which further generalizes the current nonextensive statistical mechanics. The high precision experimental and numerical data for fully developed turbulence recently discussed by Beck [23] could well be an important physical application.

Useful remarks from E. Brigatti are acknowledged, as well as partial support from PCI/MCT, CNPq, PRONEX/MCT and FAPERJ (Brazilian agencies).



## REFERENCES

- [1] C. Beck, G. S. Lewis and H. L. Swinney, Phys. Rev. E **63**, 035303 (2001).
- [2] C. Beck, Phys. Rev. Lett. **87**, 180601 (2001).
- [3] T. Arimitsu and N. Arimitsu, Physica A **305**, 218 (2002).
- [4] I. Bediaga, E.M.F. Curado and J. Miranda, Physica A **286**, 156 (2000); C. Beck, Physica A **286**, 164 (2000).
- [5] D.B. Walton and J. Rafelski, Phys. Rev. Lett. **84**, 31 (2000).
- [6] A. Upadhyaya, J.-P. Rieu, J.A. Glazier and Y. Sawada, Physica A **293**, 549 (2001).
- [7] M.L. Lyra and C. Tsallis, Phys. Rev. Lett. **80**, 53 (1998); F. Baldovin and A. Robledo, Europhys. Lett. **60**, 518 (2002); F. Baldovin and A. Robledo, Phys. Rev. E **66**, R045104 (2002); E.P. Borges, C. Tsallis, G.F.J. Ananos and P.M.C. Oliveira, Phys. Rev. Lett. **89**, 254103 (2002).
- [8] Y.S. Weinstein, S. Lloyd and C. Tsallis, Phys. Rev. Lett. **89**, 214101 (2002).
- [9] C. Anteneodo and C. Tsallis, Phys. Rev. Lett **80**, 5313 (1998); M.-C. Firpo and S. Ruffo, J. Phys. A **34**, L511 (2001); V. Latora, A. Rapisarda and C. Tsallis, Phys. Rev. E **64**, 056134 (2001); A. Campa, A. Giansanti and D. Moroni, Physica A **305**, 137 (2002); B.J.C. Cabral and C. Tsallis, Phys. Rev. E **66**, 065101(R) (2002); C. Anteneodo and R.O. Vallejos, Phys. Rev. E **65**, 016210 (2002); R.O. Vallejos and C. Anteneodo, Phys. Rev. E **66**, 021110 (2002); M.A. Montemurro, F. Tamarit and C. Anteneodo, Phys. Rev. E (2003), in press [cond-mat/0205355].
- [10] L. Borland, Phys. Rev. Lett. **89**, 098701 (2002).
- [11] S.R.A. Salinas and C. Tsallis, eds., *Nonextensive Statistical Mechanics and Thermodynamics*, Braz. J. Phys. **29**, No. 1 (1999); S. Abe and Y. Okamoto, eds., *Nonextensive Statistical Mechanics and its Applications*, Series *Lecture Notes in Physics* (Springer-

- Verlag, Berlin, 2001); G. Kaniadakis, M. Lissia and A. Rapisarda, eds., *Non Extensive Statistical Mechanics and Physical Applications*, Physica A **305**, No 1/2 (Elsevier, Amsterdam, 2002); M. Gell-Mann and C. Tsallis, eds., *Nonextensive Entropy - Interdisciplinary Applications* (Oxford University Press, 2003), in preparation; H.L. Swinney and C. Tsallis, eds., *Anomalous Distributions, Nonlinear Dynamics, and Nonextensivity*, Physica D (2003), in preparation. A regularly updated bibliography can be found at the web site <http://tsallis.cat.cbpf.br/biblio.htm> .
- [12] C. Tsallis, J. Stat. Phys. **52**, 479 (1988).
  - [13] C. Tsallis, R.S. Mendes and A.R. Plastino, Physica A **261**, 534 (1998); see also E.M.F. Curado and C. Tsallis, J. Phys. A **24**, L69 (1991) [Corrigenda: **24**, 3187 (1991) and **25**, 1019 (1992)].
  - [14] B. Lesche, J. Stat. Phys. **27**, 419 (1982).
  - [15] S. Abe, Phys. Rev. E **66**, 046134 (2002).
  - [16] R.J.V. dos Santos, J. Math. Phys. **38**, 4104 (1997); S. Abe, Phys. Lett. A **271**, 74 (2000).
  - [17] G. Wilk and Z. Włodarczyk, Phys. Rev. Lett. **84**, 2770 (2000).
  - [18] C. Beck and E.G.D. Cohen, Physica A **321** (2003), in press [cond-mat/0205097].
  - [19] C. Tsallis and A.M.C. Souza, Phys. Rev. E **67**, 0261XX (1 Feb 2003), in press [cond-mat/0206044].
  - [20] C. Beck and F. Schlogl, *Thermodynamics of Chaotic Systems* (Cambridge University Press, Cambridge, 1993).
  - [21] S. Abe, Phys. Rev. E **66**, 046134 (2002).
  - [22] J. Naudts, math-ph/0208038.

[23] C. Beck, cond-mat/0212566 (2002).